

# Northumbria Research Link

Citation: Tallent, Jamie, Goodall, Stuart, Kidgell, Dawson, Durbaba, Rade and Howatson, Glyn (2019) Compound Maximal Motor Unit Response is modulated by contraction intensity, but not contraction type in Tibialis Anterior. *Physiological Reports*, 7 (17). e14201. ISSN 2051-817X

Published by: Wiley-Blackwell

URL: <https://doi.org/10.14814/phy2.14201> <<https://doi.org/10.14814/phy2.14201>>

This version was downloaded from Northumbria Research Link:  
<http://nrl.northumbria.ac.uk/id/eprint/40030/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)





**Northumbria  
University**  
NEWCASTLE



**UniversityLibrary**

## ORIGINAL RESEARCH

# Compound maximal motor unit response is modulated by contraction intensity, but not contraction type in tibialis anterior

Jamie Tallent<sup>1,2</sup> , Stuart Goodall<sup>2</sup>, Dawson J. Kidgell<sup>3</sup>, Rade Durbaba<sup>2</sup> & Glyn Howatson<sup>2,4</sup> <sup>1</sup> School of Sport Health and Applied Science, St Mary's University, Twickenham, United Kingdom<sup>2</sup> Faculty of Health and Life Sciences, Northumbria University, Newcastle-upon-Tyne, United Kingdom<sup>3</sup> Department of Physiotherapy, School of Primary Health Care, Faculty of Medicine, Nursing and Health Sciences, Monash University, Melbourne, Australia<sup>4</sup> Water Research Group, School of Biological Sciences, North West University, Potchefstroom, South Africa**Keywords**Eccentric contractions, electromyography,  $M_{MAX}$ , peripheral nerve stimulation.**Correspondence**

Glyn Howatson, Faculty of Health and Life Sciences, Northumbria University, Newcastle-upon-Tyne NE1 8ST, United Kingdom.

Tel: +44 191 227 3575

Fax: +44 191 227 4713

E-mail: glyn.howatson@northumbria.ac.uk

**Funding Information**

No funding was received for this study.

Received: 12 July 2019; Accepted: 14 July 2019

doi: 10.14814/phy2.14201

*Physiol Rep*, 7 (17), 2019, e14201,  
<https://doi.org/10.14814/phy2.14201>**Abstract**

Determining a single compound maximal motor response ( $M_{MAX}$ ) or an average superimposed  $M_{MAX}$  response ( $M_{SUP}$ ) are commonly used reference values in experiments eliciting raw electromyographic, motor evoked potentials, H-reflexes, and V-waves. However, existing literature is limited in detailing the most appropriate method to normalize these electrophysiological measures. Due to the accessibility of assessment from a cortical and spinal perspective, the tibialis anterior is increasingly used in literature and hence investigated in this study. The aims of the present study were to examine the differences and level of agreement in  $M_{MAX}/M_{SUP}$  under different muscle actions and contraction intensities. Following a familiarization session, 22 males visited the laboratory on a single occasion.  $M_{MAX}$  was recorded under 10% isometric and 25% and 100% shortening and lengthening maximal voluntary contractions (MVC) at an angular velocity of  $15^\circ \text{ sec}^{-1}$ .  $M_{SUP}$  was also recorded during 100% shortening and lengthening with an average of five responses recorded. There were no differences in  $M_{MAX}$  or  $M_{SUP}$  between contraction types. All variables showed large, positive correlations ( $P < 0.001$ ,  $r^2 \geq 0.64$ ).  $M_{MAX}$  amplitude was larger ( $P < 0.001$ ) at 100% shortening and lengthening intensity compared to  $M_{MAX}$  amplitude at 10% isometric and 25% lengthening MVC. Bland-Altman plots revealed a bias toward higher  $M_{MAX}$  at the higher contraction intensities. Despite  $M_{SUP}$  being significantly smaller than  $M_{MAX}$  ( $P < 0.001$ ) at 100% MVC,  $M_{SUP}$  showed a large positive correlation ( $P < 0.001$ ,  $r^2 \geq 0.64$ ) with all variables. It is our recommendation that  $M_{MAX}$  should be recorded at specific contraction intensity but not necessarily a specific contraction type.

**Introduction**

Electromyographic (EMG) signals are affected by numerous factors such as preparation of the skin, electrode placement, fiber type and orientation (De Luca 1997). It is therefore critical that the EMG signal is normalized to a reference value so that data can be interpreted meaningfully. Applying supramaximal electrical stimulation to a peripheral nerve causes synchronous activation of the

muscle fibers and is known as the maximal motor unit response ( $M_{MAX}$ ; Lee and Carroll 2005). Investigations using peripherally evoked measures such as the Hoffman-reflex (H-reflex) and V-wave, along with cortically evoked measures such as motor evoked potentials (MEP), lateral spread MEP, and cervicomedullary MEP commonly use  $M_{MAX}$  as a reference value (Aagaard et al. 2002; Yamashita et al. 2002; Kidgell and Pearce 2010; Tallent et al. 2012a). These spinal and corticospinal measures have

been investigated under a variety of conditions such as changing muscle lengths, at rest and during submaximal and maximal contractions (Arányi et al. 1998; Goodall et al. 2009; Howatson et al. 2011; Tallent et al. 2012a). Understanding how  $M_{\text{MAX}}$  is modulated in different muscles, contraction intensities and types is vital in ensuring that EMG is presented in the most appropriate manner.

The  $M_{\text{MAX}}$  amplitude has been shown to increase with increasing contraction intensity in the tibialis anterior (TA: Nagata and Christianson 1995; Frigon et al. 2007) and soleus (Frigon et al. 2007), but remain unchanged in quadriceps muscles (Linnao et al. 2001a) and in the flexor carpi radialis (Lee and Carroll 2005), or even decreases in the quadriceps (Linnao et al. 2001b). In addition,  $M_{\text{MAX}}$  has been shown to increase (Gerilovsky et al. 1977; Gerilovsky et al. 1989; Frigon et al. 2007) and decrease when recorded at longer muscle lengths (Marsh et al. 1981; Kim et al. 2005; Lee and Carroll 2005). Furthermore there are conflicting findings in TA with regards to how  $M_{\text{MAX}}$  alters with changing length (Marsh et al. 1981; Frigon et al. 2007). Higher contraction intensities will cause the muscle to shorten and the tendon to become more compliant (Griffiths, 1991). A reduction in muscle length has been shown to cause an increase in synchronization and consequently an increase in  $M_{\text{MAX}}$  (Kim et al. 2005). Alternatively, phase cancellation of EMG will increase with increasing contraction intensity and may mute the response in the muscle (Keenan et al. 2006; Farina et al. 2008). Therefore, understanding  $M_{\text{MAX}}$  response at differing contraction intensities is essential from a clinical and research perspective.

Changes in muscle length might also influence  $M_{\text{MAX}}$  values during shortening and lengthening contractions. It has been recommended (Zehr, 2002) that  $M_{\text{MAX}}$  is expressed relative to the specific muscle action (i.e., shortening or lengthening muscle actions). However, evidence has shown no difference between  $M_{\text{MAX}}$  amplitude when recorded during shortening and lengthening actions (Linnao et al. 2001b; Ducloy and Martin 2005). Ensuring the reference values are recorded to a standardized muscle length appears essential in the interpretation of EMG signals.

V-wave reflects the efferent neural output during voluntary muscle activation (Aagaard et al. 2002). In the literature, V-wave is expressed relative to a mean M-wave ( $M_{\text{SUP}}$ ) during a number of maximal contractions, (Aagaard et al. 2002; Ducloy and Martin 2005; Gondin et al. 2006) or maximal peak-to-peak  $M_{\text{MAX}}$  amplitude from the same number of responses (Tallent et al. 2012b, 2013). Due to the increased potential for phase cancellation at higher contraction intensities it is unclear how these different reference values ( $M_{\text{MAX}}$  or  $M_{\text{SUP}}$ ) affect the outcome and interpretation of the V-wave. Investigating how  $M_{\text{MAX}}$  is modulated under different muscle

actions, and at varying contraction intensities might provide helpful methodological evidence for the use of  $M_{\text{MAX}}$  in experimental paradigms where neurophysiological parameters require normalization.

Therefore, the aim of this study was to investigate changes in  $M_{\text{MAX}}$  under a variety of contraction modes and intensities and examine  $M_{\text{SUP}}$  during maximal shortening and lengthening contractions in the TA. The results from this study will provide guidance for researchers in the use of  $M_{\text{MAX}}$  as a reference value.

## Methods

### Participants

Based on previous work (Kim et al. 2005) examining greater  $M_{\text{MAX}}$  amplitudes during higher contraction intensities (12%; Cohen's  $d = 0.45$ ), a total of 22 participants were recruited for the study to achieve a statistical power of 0.8 with an alpha level of 0.05. Following institutional (Northumbria University) ethical approval, 22 males (mean  $\pm$  SD, age  $23 \pm 3$  years, stature  $178.0 \pm 7.0$  cm, mass  $83.1 \pm 9.3$  kg) volunteered to participate. After being fully briefed on the experimental protocol and screened for contraindications to the procedures, volunteers provided written informed consent.

### General procedure

Two identical trials were completed, on two consecutive days at the same time of day, with the first trial used to familiarize the participants with the procedures as based on previous recommendations by our laboratory (Tallent et al. 2012a). All contractions were performed on an isokinetic dynamometer (Cybex Norm, Cybex International, NY) that was set up for ankle dorsiflexion of the dominant limb. Footedness was assessed using the questionnaire from Hebbal and Mysorekar (2006). The foot was strapped into an ankle adaptor and the knee was secured into a thigh stabilizer to prevent any extraneous movements. The hip, knee, and ankle were set at joint angles of 90, 120, and 90°, respectively, according to the manufacturer's instructions. Shortening and lengthening contractions consisted of participants moving through a range of 30° ( $\pm 15^\circ$  from the ankle at 90°) at an angular velocity of  $15^\circ \cdot \text{sec}^{-1}$ . Shortening and lengthening contractions began at an ankle angle of 105° and 75° respectively. For shortening muscle actions, participants were instructed to assist the movement of the foot adaptor, and for lengthening the actions required participants to resist movement of the foot adaptor. All responses (torque and EMG) were recorded as the ankle joint passed through anatomical zero (90°). To ensure torque and

EMG were recorded at the correct angle, a trigger was set to automatically sweep as the ankle passed 90°. Once secured in the isokinetic dynamometer, participants initially performed shortening, lengthening and isometric MVCs. The highest torque in each muscle action (shortening, lengthening, and isometric) from three trials was recorded as the contraction-specific MVC.

The  $M_{\text{MAX}}$  was recorded at 10% of isometric, 25% and 100% shortening and lengthening MVC. A 10% isometric contraction is often used to stabilize the H-reflex in the TA (Griffin & Cafarelli, 2007; Tallent et al. 2012a), and consequently this was considered the resting  $M_{\text{MAX}}$  value. The stimulation intensity for eliciting  $M_{\text{MAX}}$  was set at 150% above a plateau in peak-to-peak  $M_{\text{MAX}}$  amplitude. This was recorded through an increasing stimulation intensity at 10% isometric MVC and verified during 25% shortening and lengthening contractions. Establishing  $M_{\text{MAX}}$  took around 64 gradually increasing intensity pulses at 10% isometric MVC.  $M_{\text{SUP}}$  was calculated from the average of 5 traces at 100% shortening and lengthening MVC, whilst  $M_{\text{MAX}}$  during a maximal contraction was recorded as the greatest peak-to-peak amplitude of the 5 contractions. The order of contraction intensity (10%, 25%, 100%) and type (shortening and lengthening) was randomized.

### Percutaneous nerve stimulation

Searching for optimal site of stimulation began below the head of the fibula, over the peroneal nerve. A 1 msec electrical stimulation was administered using a 40 mm diameter cathode/anode arrangement (Digitimer DS7AH, Welwyn Garden City, Hertfordshire, UK). Once the optimal site was located, the sight was marked with semi-permanent ink. The cathode/anode was strapped to the participants' leg for the entirety of the experiment.

### EMG

Bipolar surface EMG was recorded over the TA using electrodes (22 mm diameter, model; Kendall, Tyco Healthcare Group, Mansfield, MA) spaced 2 cm apart. The reference electrode was placed over the medial malleolus, whilst the TA electrodes were placed at one-third distance of the line between the tip of the fibula and the tip of the medial malleolus (Hermens et al. 2000). All sites were shaved, abraded, and then wiped clean with an alcohol swab prior to electrode placement. The EMG was amplified ( $\times 1000$ ), band pass filtered (10–1000 Hz), and sampled at 5 kHz (CED Power 1401, Cambridge Electronic Design, Cambridge, UK). M-waves were recorded during a 500 msec window, starting 50 msec before anatomical zero. Once  $M_{\text{MAX}}$  stimulator was established, all further analyses were performed off-line.

### Torque

To ensure that participants reached the required target torque level, real time feedback was provided on a computer monitor positioned 1 m away. Live feedback was displayed on the monitor of the dynamometer to provide feedback on target forces to achieve during each condition. The torque signal was sampled at 5 kHz, extracted from the dynamometer and synchronized with the EMG signal and analysed off line (Signal v3.0, Cambridge Electronics, Cambridge, UK).

### Statistics

A one-way ANOVA was used to detect differences between  $M_{\text{MAX}}$  at 10% isometric MVC, 25%, 100% and  $M_{\text{SUP}}$  at 100% shortening and lengthening MVC. Where necessary, LSD post-hoc analysis was used to make pairwise comparisons with 95% CI (SPSS, v20.0, Chicago, IL). Coefficient of determination and the limits of agreement (Bland and Altman 1986) with 95% CI were also calculated between the variables (GraphPad Software Inc, La Jolla, CA). Correlation coefficients were determined as 0.0–0.1 = trivial, 0.10–0.3, small, 0.3–0.5 = moderate, 0.5–0.7 = large, and 0.7–0.9 = very large (Hopkins, 2009). Effect sizes ( $\eta^2$ ) were defined as: 0.2 trivial, 0.21–0.6 = small, 0.61–1.2 = moderate, 1.21–1.99 = large;  $>2.0$  = very large.

### Results

Isometric contractions were conducted at an average of  $8.28 \pm 3.21\%$  (target = 10%) of isometric MVC, shortening at  $26.1 \pm 3.66\%$  (target = 25%),  $95.6 \pm 11.8\%$  (target = 100%) of shortening MVC and lengthening at  $27.1 \pm 4.12\%$  (target = 25%),  $96.2 \pm 9.97\%$  (target = 100%) of lengthening MVC. There was no significant difference ( $P > 0.05$ ) between lengthening and shortening contraction intensities, showing that contractions were conducted at the same relative intensity.

Figure 1 shows individual and average  $M_{\text{MAX}}/M_{\text{SUP}}$  amplitudes during varying isometric, shortening, and lengthening contractions intensities and a representative trace. The ANOVA revealed there were significant differences in  $M_{\text{MAX}}$  amplitude between conditions ( $F_{(6)} = 6.96$ ;  $P < 0.001$ ;  $\eta^2 = 0.25$ ). Post Hoc analysis showed 10% isometric  $M_{\text{MAX}}$  MVC was significantly lower than 25% shortening  $M_{\text{MAX}}$  ( $P = 0.03$ ; 95% CI;  $-0.03$  to  $-0.69$  mV), 25% lengthening  $M_{\text{MAX}}$  ( $P = 0.03$ ; 95% CI;  $-0.05$  to  $-0.64$  mV), 100% shortening  $M_{\text{MAX}}$  ( $P < 0.01$ ; 95% CI;  $-0.40$  to  $-1.32$  mV), 100% lengthening  $M_{\text{MAX}}$  ( $P < 0.01$ ; 95% CI;  $-0.37$  to  $-1.32$  mV).  $M_{\text{MAX}}$  was significantly higher at 100% shortening ( $P = 0.02$ ; 95% CI;  $0.11$  to  $1.03$  mV) and 100%

lengthening ( $P = 0.02$ ; 95% CI; 0.08–1.03) compared to 25% lengthening  $M_{MAX}$ .


All  $M_{MAX}$  amplitudes were significantly ( $P < 0.001$ ) correlated across intensities ( $r^2, \geq 0.64$ ). The highest correlations were between contraction types at the same intensity with  $M_{MAX}$  (100% MVC  $r^2 = 0.87$ ; 25% MVC  $r^2 = 0.86$ ). Bland-Altman plots showed a bias toward higher  $M_{MAX}$  values at higher contraction intensities (Fig. 2). There was no bias between shortening and lengthening contractions. Similarly, isometric  $M_{MAX}$  and  $M_{SUP}$  showed no bias.

## Discussion


It has been reported that EMG should be normalized to  $M_{MAX}$  under the same muscle action and contraction intensity (Zehr, 2002; Duclay and Martin 2005). This study offers further insight into the influence that contraction conditions may affect the amplitude of  $M_{MAX}$

amplitude. Specifically, the main findings were, (1) there was no difference between  $M_{MAX}$  amplitudes when recorded at like-intensities during shortening and lengthening contractions; (2)  $M_{MAX}$  was influenced by intensity of the contraction, with an increase and systematic bias to an increase  $M_{MAX}$  during higher intensity contractions; and (3)  $M_{MAX}$  at 100% MVC was greater compared to  $M_{SUP}$  at 100% MVC. However,  $M_{SUP}$  was not different to  $M_{MAX}$  at 10% MVC, showing little systematic bias and was strongly correlated ( $r^2 \geq 0.64$ ).

It has been recommended that when using  $M_{MAX}$  as a reference value, it should be recorded under the same contraction intensity as the variable being investigated (Zehr, 2002). The results in this study indicated that with increased contraction intensity the peak-to-peak  $M_{MAX}$  amplitude increased. Previous work has shown similar results in contraction intensities ranging from 40 to 80% isometric MVC in TA (Nagata and Christianson 1995) and 10–30% isometric MVC in TA and the soleus (Frigon et al.



**Figure 1.** Clear dots represent individual responses at different  $M_{MAX}$  contraction intensities and contraction types (A). Bars represent mean  $M_{MAX}$  and  $M_{SUP}$  responses (mean  $\pm$  SD) (B). Representative trace from a single participant of  $M_{MAX}$  recorded at 10% ISO, SHO and LEN 25% and 100% MVC, SHO and LEN  $M_{SUP}$  (C). ISO, Isometric, SHO, Shortening, LEN, lengthening; \*denotes significantly ( $P < 0.05$ ) different from 25% and 100%, SHO and LEN MVC  $M_{MAX}$ ; \*\*denotes significantly different from 100% SHO and LEN  $M_{MAX}$ ; \*\*\*denotes significantly different from SHO and LEN  $M_{SUP}$ .



**Figure 2.** Bland-Altman plots for  $M_{MAX}$  and  $M_{SUP}$  (mV) across varying contraction intensities and type. Dashed line indicated change in mean with 95% confidence intervals. Dots represent individual responses.



2007). Although this effect is reported previously and supported by the current study, the exact mechanisms for this remain unclear. Frigon et al. (2007) suggested that  $M_{\text{MAX}}$  increased at higher contraction intensities because the muscle length has been shown to be up to 28% shorter at the same joint angle (Griffiths, 1991), and thus, could improve the synchronization of the action potential. However, contrary with our findings, other authors have reported no change (Linnaam et al. 2001a; Lee and Carroll 2005) or even a decrease (Linnaam et al. 2001b) in  $M_{\text{MAX}}$  with increasing contraction intensities. The high degree of variability between subjects might explain why the literature offers little consistency (Lee and Carroll 2005). In addition, phase cancellation has been shown to reduce the EMG response at the muscle during higher contraction intensities (Keenan et al. 2006; Farina et al. 2008). If the responses in EMG are muted at higher contraction intensities then the lack of change in  $M_{\text{MAX}}$  amplitude appears to be associated with the limitations in surface EMG recording (Farina et al. 2014).

Our results support previous findings that showed no difference in  $M_{\text{MAX}}$  under shortening and lengthening contractions (Linnaam et al. 2001b; Duclay and Martin 2005) when measured at the same joint angle. Supramaximal stimulation of the peripheral nerve at shorter muscle lengths improves the synchronization of the action potential (Kim et al. 2005). With an enhanced synchronization of the action potential there is an increase in  $M_{\text{MAX}}$  (Kim et al. 2005). However, the varying pennation angle of muscles might explain why not all studies have found increases in  $M_{\text{MAX}}$  at shorter muscle lengths (Gerilovsky et al. 1977; Gerilovsky et al. 1989; Frigon et al. 2007). Furthermore, it is expected that an increase in  $M_{\text{MAX}}$  at shorter muscle lengths should be associated with decreased duration of  $M_{\text{MAX}}$ , although this is not consistently observed (Frigon et al. 2007). In our study,  $M_{\text{MAX}}$  was recorded during shortening and lengthening muscle contractions and importantly, electrical stimulation was delivered at the same joint angle, with the assumption that the muscle was at the same length. Furthermore, current data also showed a strong positive correlation, and a good level of agreement, between  $M_{\text{MAX}}$  during shortening and lengthening muscle actions. Thus, it appears that EMG signals do not necessarily need to be expressed relative to a contraction specific  $M_{\text{MAX}}$ , rather, the joint angle should be consistent (Nagata and Christianson 1995; Kim et al. 2005; Frigon et al. 2007). A high level of agreement and a strong correlation was found between shortening and lengthening muscle actions, despite lengthening muscle actions generating a higher level of absolute torque. The differences in  $M_{\text{MAX}}$  at an 'absolute' torque might explain why there is a small discrepancy between shortening and lengthening  $M_{\text{MAX}}$  at the same relative intensity.

Unlike MEP's, H-reflex, and EMG signals, V-wave is expressed relative to an  $M_{\text{SUP}}$  (Aagaard et al. 2002; Duclay and Martin 2005; Gondin et al. 2006) or  $M_{\text{MAX}}$  (Tallent et al. 2012b; 2013). In this study, there was no difference in  $M_{\text{MAX}}$  at a low intensity contraction ( $\leq 25\%$ ) and  $M_{\text{SUP}}$ . This would suggest that EMG/V-waves recorded during an MVC could be expressed relative to a low intensity  $M_{\text{MAX}}$  contraction. There was also good level of agreement between  $M_{\text{SUP}}$  and  $M_{\text{MAX}}$  at low intensity contractions suggesting these values could be used interchangeably, although in the interest of rigor, it would be sensible to use a single well controlled  $M_{\text{MAX}}$  measure to normalize all conditions.

## Conclusion

The results from this study show that  $M_{\text{MAX}}$  is not altered by shortening or lengthening contraction type, but is modulated with changes in contraction intensity. Possible mechanisms may be due to the shortened muscle lengths at the higher contraction intensities.  $M_{\text{MAX}}$  should be used relative to task specific contraction intensities and it is vital that it is recorded under consistent reproducible conditions. No differences were seen between  $M_{\text{MAX}}$  at low intensity contractions and  $M_{\text{SUP}}$  at 100% MVC. There was also low systematic bias and strong correlations suggesting that V-wave can be expressed relative to  $M_{\text{MAX}}$  recorded during low intensity contractions or  $M_{\text{SUP}}$  at 100% MVC. It is our recommendation that  $M_{\text{MAX}}$  should be recorded at specific contraction intensities but not necessarily a specific contraction type. However, consistency of  $M_{\text{MAX}}$  recording throughout the experiment is vital.

## Acknowledgments

No acknowledgments.

## Conflict of Interest

The authors have no competing interests to declare, financial, or otherwise.

## References


- Aagaard, P., E. B. Simonsen, J. L. Andersen, P. Magnusson, and P. Dyhre-Poulsen. 2002. Neural adaptation to resistance training: changes in evoked V-wave and H-reflex responses. *J. Appl. Physiol.* 92:2309–2318.
- Arányi, Z., J. Mathis, C. W. Hess, and K. M. Rösler. 1998. Task-dependent facilitation of motor evoked potentials during dynamic and steady muscle contractions. *Muscle Nerve* 21:1309–1316.

- Bland, J. M., and D. G. Altman. 1986. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* 1:307–310.
- Duclay, J., and A. Martin. 2005. Evoked H-reflex and V-Wave responses during maximal isometric, concentric, and eccentric muscle contraction. *J. Neurophysiol.* 94:3555–3562.
- Farina, D., C. Cescon, F. Negro, and R. M. Enoka. 2008. Amplitude cancellation of motor-unit action potentials in the surface electromyogram can be estimated with spike-triggered averaging. *J. Neurophysiol.* 100:431–440.
- Farina, D., R. Merletti, and R. M. Enoka. 2014. The extraction of neural strategies from the surface EMG: an update. *J. Appl. Physiol.* (1985) 117:1215–1230.
- Frigon, A., T. J. Carroll, K. E. Jones, E. P. Zehr, and D. F. Collins. 2007. Ankle position and voluntary contraction alter maximal M waves in soleus and tibialis anterior. *Muscle Nerve* 35:756–766.
- Gerilovsky, L., A. Gydikov, and N. Radicheva. 1977. Changes in the shape of the extraterritorial potentials of tonic motor units, M- and H-responses of triceps surae muscles at different muscle lengths and under conditions of voluntary activation. *Exp. Neurol.* 56:91–101.
- Gerilovsky, L., P. Tsvetinov, and G. Trenkova. 1989. Peripheral effects on the amplitude of monopolar and bipolar H-reflex potentials from the soleus muscle. *Exp. Brain Res.* 76:173–181.
- Gondin, J., J. Duclay, and A. Martin. 2006. Soleus- and gastrocnemii-evoked V-wave responses increase after neuromuscular electrical stimulation training. *J. Neurophysiol.* 95:3328–3335.
- Goodall, S., L. M. Romer, and E. Z. Ross. 2009. Voluntary activation of human knee extensors measured using transcranial magnetic stimulation. *Exp. Physiol.* 94:995–1004.
- Griffin, L., and E. Cafarelli. 2007. Transcranial magnetic stimulation during resistance training of the tibialis anterior muscle. *J. Electromyogr. Kinesiol.* 17:446–452. <https://doi.org/10.1016/j.jelekin.2006.05.001>.
- Griffiths, R. I. 1991. Shortening of muscle fibres during stretch of the active cat medial gastrocnemius muscle: the role of tendon compliance. *J. Physiol.* 436:219–236.
- Hebbal, G. V., and V. R. Mysorekar. 2006. Evaluation of some tasks used for specifying handedness and footedness. *Percept. Mot. Skills* 102:163–164.
- Hermens, H. J., B. Freriks, C. Disselhorst-Klug, and G. Rau. 2000. Development of recommendations for SEMG sensors and sensor placement procedures. *J. Electromyogr. Kinesiol.* 10:361–374.
- Hopkins, A. (2009). A scale of magnitudes for effect statistics. A new view of statistics.
- Howatson, G., M. B. Taylor, P. Rider, B. R. Motawar, M. P. McNally, S. Solnik, et al. 2011. Ipsilateral motor cortical responses to TMS during lengthening and shortening of the contralateral wrist flexors. *Eur. J. Neurosci.* 33:978–990.
- Keenan, K. G., D. Farina, R. Merletti, and R. M. Enoka. 2006. Amplitude cancellation reduces the size of motor unit potentials averaged from the surface EMG. *J. Appl. Physiol.* (1985) 100:1928–1937.
- Kidgell, D. J., and A. J. Pearce. 2010. Corticospinal properties following short-term strength training of an intrinsic hand muscle. *Hum. Mov. Sci.* 29:631–641.
- Kim, B. J., E. S. Date, B. K. Park, B. Y. Choi, and S. H. Lee. 2005. Physiologic changes of compound muscle action potentials related to voluntary contraction and muscle length in carpal tunnel syndrome. *J. Electromyogr. Kinesiol.* 15:275–281.
- Lee, M., and T. J. Carroll. 2005. The amplitude of  $M_{MAX}$  in human wrist flexors varies during different muscle contractions despite constant posture. *J. Neurosci. Methods* 149:95–100.
- Linnamo, V., V. Strojnik, and P. V. Komi. 2001a. Electromyogram power spectrum and features of the superimposed maximal M-wave during voluntary isometric actions in humans at different activation levels. *Eur. J. Appl. Physiol.* 86:28–33.
- Linnamo, V., V. Strojnik, and P. V. Komi. 2001b. EMG power spectrum and maximal M-wave during eccentric and concentric actions at different force levels. *Acta Physiol. Pharmacol. Bulg.* 26:33–36.
- De Luca, C. J. 1997. The use of electromyography in biomechanics. *J. Appl. Biomech.* 13:18.
- Marsh, E., D. Sale, A. J. McComas, and J. Quinlan. 1981. Influence of joint position on ankle dorsiflexion in humans. *J. Appl. Physiol. Respir. Environ. Exerc. Physiol.* 51:160–167.
- Nagata, A., and J. C. Christianson. 1995. M-wave modulation at relative levels of maximal voluntary contraction. *Eur. J. Appl. Physiol. Occup. Physiol.* 71:77–86.
- Tallent, J., S. Goodall, T. Hortobágyi, A. St Clair Gibson, D. N. French, and G. Howatson. 2012a. Repeatability of corticospinal and spinal measures during lengthening and shortening contractions in the human tibialis anterior muscle. *PLoS ONE* 7:e35930.
- Tallent, J., S. Goodall, T. Hortobágyi, A. St Clair Gibson, D. N. French, and G. Howatson. 2012b. Recovery time of motor evoked potentials following lengthening and shortening muscle action in the tibialis anterior. *J. Clin. Neurosci.* 19:1328–1329. <https://doi.org/10.1016/j.jocn.2012.1001.1022>.
- Tallent, J., S. Goodall, T. Hortobágyi, A. St Clair Gibson, and G. Howatson. 2013. Corticospinal responses of resistance-trained and un-trained males during dynamic muscle contractions. *J. Electromyogr. Kinesiol.* 23:1075–1081.
- Yamashita, S., T. Kawaguchi, M. Fukuda, K. Suzuki, M. Watanabe, R. Tanaka, et al. 2002. Lateral spread response elicited by double stimulation in patients with hemifacial spasm. *Muscle Nerve* 25:845–849.
- Zehr, P. 2002. Considerations for use of the Hoffmann reflex in exercise studies. *Eur. J. Appl. Physiol.* 86:455–468.



# Graphical Abstract

The contents of this page will be used as part of the graphical abstract of html only. It will not be published as part of main.



The study investigated the effect of contraction type and intensity on maximal compound action potential in the tibialis anterior. It is our recommendation that maximal compound action potential should be recorded at specific contraction intensity but not necessarily a specific contraction type.